Attention control comparisons with SLT for people with aphasia following stroke: methodological concerns raised following a systematic review

Brady, Marian C.; Godwin, Jon; Kelly, Helen; Enderby, Pam; Elders, Andrew; Campbell, Pauline

Published in:
Clinical Rehabilitation

DOI:
[10.1177/0269215518780487](https://doi.org/10.1177/0269215518780487)

Publication date:
2018

Document Version
Author accepted manuscript

[Link to publication in ResearchOnline](#)

Citation for published version (Harvard):

Brady, MC, Godwin, J, Kelly, H, Enderby, P, Elders, A & Campbell, P 2018, 'Attention control comparisons with SLT for people with aphasia following stroke: methodological concerns raised following a systematic review', *Clinical Rehabilitation*, vol. 32, no. 10, pp. 1383–1395. <https://doi.org/10.1177/0269215518780487>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please view our takedown policy at <https://edshare.gcu.ac.uk/id/eprint/5179> for details of how to contact us.

ABSTRACT

Objective

Attention control comparisons in trials of stroke rehabilitation require care to minimise the risk of comparison choice bias. We compared the similarities and differences in speech and language therapy and social support control interventions for people with aphasia.

Data Sources: Trial data from the 2016 Cochrane systematic review of speech and language therapy for aphasia after stroke

Methods: Direct and indirect comparisons between speech and language therapy, social support and no-therapy controls. We double-data extracted intervention details using the template for intervention description and replication. Standardized mean differences and risk ratios (95% confidence intervals) were calculated.

Results: Seven trials compared speech and language therapy with social support (n=447). Interventions were matched in format, frequency, intensity, duration and dose. Procedures and materials were often shared across interventions. Social support providers received specialist training and support. Targeted language rehabilitation was only described in therapy interventions. Higher dropout ($P = 0.005$, OR 0.51 95% CI 0.32 to 0.81) and non-adherence to social support interventions ($P < 0.00001$, OR 0.18, 95% CI 0.09 to 0.37) indicates imbalanced completion rates increasing the risk of control comparison bias.

Conclusion:

Distinctions between social support and therapy interventions were eroded. Theoretically based language rehabilitation was the remaining difference in therapy interventions. Social support is an important formal language rehabilitation adjunct. Therapists should continue to enable those close to the person with aphasia to provide tailored communication support, functional language stimulation and opportunities to apply rehabilitation gains. Systematic group differences in completion rates is a design-related risk of bias in outcomes observed.

Attention control comparisons with speech and language therapy for people with aphasia following stroke: methodological concerns raised following a systematic review

Introduction:

Rehabilitation of aphasia (the impairment of language after stroke or other neurological disorder), seeks to maximise an individual's return to communication activities and participation. In a recent Cochrane review ¹ speech and language therapy interventions were found to benefit people with aphasia as determined by their performance on measures of functional communication and language impairment (language expression, reading and writing), when compared to people with no access to such therapy ¹. There was no clear evidence in the context of comparisons with social support attention control interventions.

Rigorous evaluation of rehabilitation interventions should be conducted in the context of a randomised comparison which might include randomisation to a usual care group or an attention control group. Suitable control comparator groups ensure that intervention effects can be evaluated in isolation from other factors which might influence outcome; such as natural recovery, increased healthcare professional attention, benefits from trial participation and trial expectations ². For example, participants in stroke research have been found to receive better care than non-participating peers ³. In the context of trials of speech and language therapy for aphasia after stroke, social support attention control comparisons have been advocated ⁴ given that no attention may be questionable on ethical grounds.

People with aphasia are at high risk of social isolation ^{34, 35}. Social support is likely to benefit their wellbeing and quality of life which in turn could benefit their engagement with rehabilitation and

social participation. Clinical psychology and psychotherapy researchers have highlighted the importance of social support, encouragement and a therapeutic relationship as active components of effective therapeutic interventions^{5,6}. Regular social support shares some characteristics and benefits with specific therapeutic interventions. Speech and language therapy, incorporating conversational practice, for people with aphasia is one such intervention. Social support has been found to be an unsuitable control comparator in evaluations of the effectiveness of communication based therapeutic interventions^{5,6}.

Regular participation in social situations provides scheduled opportunities for practicing functionally relevant language use, a key outcome for most aphasia rehabilitation interventions. As social support interventions are inherently language based such attention control interventions are not easily distinguished from more specific speech and language therapy interventions.

Methodologically, it is important to preserve a clear distinction between trial group interventions; otherwise a trial may risk underestimating the effectiveness of an experimental intervention. Trials of other stroke rehabilitation interventions preserve this distinction. For example, in a recent review of physical rehabilitation interventions to improve function and mobility after stroke 12 of 96 trials included an attention control group⁷. The content of those control interventions ranged from upper limb therapy (n=6 trials), cognitive training (n=4 trials), massage (n=1) and a socially based educational group (n=1)⁷. None offered an attention control intervention that facilitated the functional application or practice opportunities for the target activity, such as a volunteer-supported walk in the park, participation in a walking group or similar.

Our recent systematic review of speech and language therapy interventions for people with aphasia identified 57 randomised controlled trials that compared a speech and language intervention with

another speech and language therapy intervention (36 trials; n= 1242) with no access to therapy (22 trials; n=1620) or with social support (7 trials; n=447) described by the trialists as an attention control ¹. In this paper we aim to (i) examine the similarities and differences in the speech and language therapy and social support attention control interventions compared within these randomised controlled trials; (ii) present the evidence on the comparative effectiveness of speech and language therapy and social support interventions on language outcomes, intervention adherence and trial dropouts using meta-analyses and indirect comparisons; (iii) consider the relative acceptability and potential risk of bias in the use of social support interventions in trials of the clinical effectiveness of speech and language therapy for aphasia after stroke.

Methods

This analysis starts from a systematic review and meta-analysis of randomised controlled trials that evaluated speech and language therapy interventions designed to improve language or communication amongst adults with aphasia as a result of stroke and which were compared to social support interventions. Our review was conducted to agreed methodological and reporting standards ^{8,9}, and is reported in detail elsewhere. ¹ Briefly we systematically searched several electronic databases including **Central** Register of Controlled Trials, MEDLINE, EMBASE, Cumulative Index to Nursing and Allied Health Literature, Allied and Complementary Medicine Database.

Speech and language therapy interventions were defined as any form of targeted practice, tasks or methodologies which had the aim of improving language or communication abilities of people with aphasia, regardless of the individual delivering the intervention. Social support interventions were defined as stimulating functionally relevant social language use

in a naturalistic setting following an intervention regimen or schedule detailed within the trial protocol and often informed by a speech and language therapist's assessment or intervention manual but which did not include components that targeted specific aspects of language rehabilitation. Descriptions of complex non-pharmacological interventions are known to be insufficient ¹⁰.

For this analysis, we extracted available information on the interventions from published papers, and we contacted the primary research teams to supplement data extraction. We used the Template for Intervention Description and Replication (TIDieR) to support systematic data extraction ¹⁰. We then profiled the interventions using these headings to consider the similarity and differences between speech and language therapy and social support interventions. Data extraction was conducted independently by two review authors with a third resolving any disputes.

We conducted direct and indirect comparisons between speech and language therapy interventions with social support comparators and with no therapy controls. Outcomes of relevance included functional communication, receptive language, expressive language or aphasia severity. We also considered the number of participants that dropped out from the trials (during the interventions or at the outcome assessment time points and for any reason) and the extent of non-adherence to allocated interventions ⁸. Where suitable statistical summary data were available we combined the selected outcome data in pooled meta-analyses and indirect comparisons. Where a single outcome measure was assessed across trials which used different measurement tools (thus producing indirectly comparable data) we were unable to assume a common treatment effect and combined the data using

standardised mean differences. For binary outcomes (attrition data) we combined the data using relative risk ratios.

Pooled effect sizes for speech and language therapy versus no speech and language therapy and speech and language therapy versus social support were calculated from random effects meta-analyses using the DerSimonian and Laird method ¹¹. Effect sizes for social support versus no SLT were then estimated based on Bucher's method ¹² for adjusted indirect comparisons. In these analyses, standardised mean differences (with Hedge's adjusted g to correct small sample bias ¹³) were used for aphasia outcomes. Risk ratios were used to compare dropout and non-adherence rates. Where randomised participants were at risk of being included twice in a single meta-analysis we split the number of participants in the shared group across the two trials ⁸. For continuous data, the mean and standard deviation values remained the same. For dichotomous data, we split both the number of events and total number of patients. We assessed heterogeneity using the I² statistic, where values of greater than 50% indicated substantial heterogeneity. To address potential heterogeneity, we used random-effects models to pool the data and the source of any substantial heterogeneity was investigated.

Results

Intervention description – speech and language therapy versus social support

We included seven trials (n=447 randomised participants) that compared speech and language therapy with a social support intervention ^{14a-b, 15-19, 20a-b}. Two were three armed trials thus there were nine randomised comparisons. ^{14a-b, 20a-b} (Table 1). One was a cross-

over trial where we extracted data up to the point of intervention cross-over¹⁶. We found no evidence of a difference between the groups as measured on functional (activity and participation) or language outcome measures. However, more participants dropped out (n=65 for any reason) or failed to adhere (n=45) to the social support interventions than those that were allocated to speech and language therapy (n=40 dropouts p=0.005; n=11 non-adherence p<0.00001). Thus social support interventions were maybe less acceptable to the participants than targeted language intervention³⁷. Below we consider the two interventions on the data items from the TIDieR checklist¹⁰ (Tables 2 and 3).

Why

Social support interventions were almost exclusively described by trialists as an intervention to control for the effects of social contact, encouragement to communicate and attention within the trial (“attention control”^{14a-b, 19}, control for “attention effect”¹⁵ or “control the effects of social contact”¹⁷, “control sessions”¹⁸). One described active ingredients within the intervention which was “stimulation-orientated, designed to provide psychological support and work on communication in unstructured settings”^{20a-b}.

In contrast, where described, the speech and language therapy interventions sought to facilitate language recovery^{14a-b, 16-20}.

What

Most social support interventions aimed to encourage and stimulate conversation facilitated by the intervention provider. One trial specifically encouraged a participant-led

social interaction¹⁹ where participants regularly took part in a local class or group activity of their choice which was not detailed in the trial report¹⁷. Three described providing psychological support or building a rapport with participants^{19, 20a-b}. Those providing social support had access to a manual of suitable conversation topics^{16, 19}, a conversational support strategy handbook^{14a-b}, information on participants' aphasia^{15, 20a-b}, assessment scores and support requirements¹⁵. One trial employed a formal narrative re-telling task¹⁸ while social support participation in other trials involved the creative arts^{17, 19}, listening to music, watching television, reading, playing approved board games or gardening¹⁹. The specific materials were rarely reported.

In contrast speech and language therapy interventions were usually defined and detailed in the protocol^{14, 17-20a-b}. Where described, the therapy included targeted stimulation of specific language structures and skills¹⁹ including comprehension^{16, 17, 20a-b} and expressive skills^{14, 16-18, 20a-b} including reading and writing^{16, 20a}. Two speech and language therapy interventions were left to the therapists' discretion^{15, 16}. Intervention fidelity monitoring was described in six trials^{19, 20a-b} where three monitored a percentage of the overall sessions^{14a-b, 15}.

Who

Where reported, social support was provided by volunteers, psychologists, nurses, researchers, community-based facilitators or speech and language therapy students (Table 2). In six trials the social support providers were trained in the delivery of the intervention^{19, 15, 16, 18, 20a-b14a-b}, had information on the participants' aphasia^{15, 20a-b}, their formal aphasia assessment scores and

communication support needs¹⁵, a manual supporting the intervention^{14a-b, 15, 16, 19} and ongoing support from the SLT research team in initiation and adaptation of communication^{15, 14a-b}.

Participants across groups were similar except for one trial¹⁵ where those that received speech and language therapy were significantly older than those that received social support.

Speech and language therapy was typically provided by professionally qualified speech and language therapists except for in two trials, one where it was delivered by a trained researcher,¹⁸ and one where therapy delivered by the therapist was augmented by additional input from a family member¹⁷.

How

The model of intervention delivery was similar. Most social support and therapy interventions were provided on a one-to-one and face-to-face basis. In two cases both interventions were provided at group level¹⁷ or via a computer interface^{14a-b}.

Where

Social support and formal therapy interventions were, where reported, usually provided in similar settings. One trial provided social support in 'unstructured settings'^{20a-b} while the location of the comparison therapy group was unreported. Another provided social support at home while formal therapy was provided in clinic with home practice^{14a-b}.

When and how much

Most intervention comparisons were matched for frequency (sessions weekly), intensity (hours of intervention weekly), duration (overall length of intervention) and dose (total

hours of intervention delivered) (Table 3). Interventions were provided during 2 to 4 sessions, for 2 to 3 hours weekly, over one to 12 months. Intervention dose ranged from 8 to 156 hours. In one trial the social support weekly (minimum of 3 hours) and total dose (52 hours) of intervention was less than the speech and language therapy comparison (5 hours and up to 160 hours)¹⁷ (Table 2). In another it was difficult to compare weekly frequency and intensity but the average dose and total duration of social support (15 hours over 16 weeks) was similar to speech and language therapy provision (18 hours over 16 weeks)¹⁹.

Effects of Interventions

Statistical data which permitted inclusion within speech and language versus social support meta-analyses were available for six trials^{14a-b, 15, 16, 18, 19} (Table 4). Suitable outcome data were unavailable for the remaining trials^{17, 20a-b}. In three instances where data permitted pooled meta-analysis, there was no evidence of a difference between the groups' performance on functional communication or auditory comprehension. One small trial (n=18)¹⁶ reported that the participants that received social support gained significant benefit on measures of general receptive language, expressive language, writing, word fluency and aphasia severity (Table 4).

Based on data from five trials significantly more participants receiving social support dropped out of the trials during or at post-intervention assessments compared to the number lost to speech and language therapy interventions (p=0.012, risk ratio 0.65 95% confidence interval 0.46 to 0.91) (Table 5). None of the five trials employed an intention to treat analysis approach. An additional 13 people were excluded from the primary trials for failing to complete intervention protocols but their allocation was unclear and so they were not included in our analysis¹⁶. When we considered voluntary withdrawal (non-adherence to the intervention) the difference became more pronounced

^{15, 17, 19, 20a-b} ($p < 0.001$, risk ratio 0.24, 95% confidence interval 0.12 to 0.47) (Table 5). Four additional participants voluntarily withdrew from the social support group because of 'volunteer problems' ¹⁵.

Using the group summary data from the recently updated Cochrane review of speech and language therapy for aphasia after stroke ¹ we conducted direct and (for the first time) indirect comparisons using the language outcome data (functional communication, receptive and expressive speech and severity of aphasia). Our previous meta-analyses found that speech and language therapy was significantly more effective than no SLT ¹. We found some very weak evidence (based on one small trial $n=18$) that participants receiving social support performed significantly better on some language outcomes (writing, general receptive, expressive language and severity) than participants that received speech and language therapy (Table 4).

The novel indirect comparisons presented within this paper found limited evidence to suggest that participants receiving social support interventions performed better on measures of writing, general expressive language and severity of aphasia compared to participants that received no therapy (Table 4). Heterogeneity was low. Similarly we found no evidence of a difference in dropout and non-adherence rates between social support groups and no therapy control groups. However, this may be due to the small sample sizes of included studies (Table 5).

Discussion

While speech and language therapy benefits people with aphasia compared to no intervention, our meta-analysis found no evidence of an impact on participant outcomes when direct SLT intervention was compared in a small number of trials (involving few participants) to a social support attention control. The significantly higher dropout (differential dropout) and non-adherence rates amongst

participants randomly allocated to social support interventions compared to speech and language therapy is a potential source of bias which threatens the validity of the results and is a recognised source of concern^{33, 36}. Those participants that remained in the trial and completed the outcome assessments may have differed from those that dropped out. Evidence of such differences between the interventions raises questions about the acceptability and suitability of social support as a control comparison in randomised controlled trials of aphasia therapy^{33, 37}.

We found that the format, location, frequency and intensity (weekly) and overall duration and dose of social support and speech and language therapy interventions were matched across comparisons. Only one of seven trials ensured a differential dose between the comparisons¹⁷. Social support providers received training by a speech and language therapist prior to providing that support. . Therapists also provided information on the patient's abilities and difficulties following assessment, training, support and materials to the providers of social support. Interventional procedures and materials developed and supplied by the therapy research team were also shared across interventions in some trials

In contrast, targeted rehabilitation of specific language structures (informed by patient preference or assessment findings) and a clear language recovery goal were only described in speech and language interventions. The extent of the similarities observed between the speech and language therapy and social support interventions however have diluted any distinction between targeted rehabilitation activities and a planned attention control intervention. Indirect comparisons highlighted some significant differences in outcome measures between participants in social support activities and those that received no speech and language therapy, but the data was generally based on small sample sizes and a small number of trials.

Previous systematic reviews in this field have tended to focus on the effectiveness of therapy interventions with limited attention given to the processes followed in the comparator groups²⁸⁻³¹. Our systematic consideration of both the experimental therapy intervention and the social support attention control comparisons highlighted important clinical and methodological issues.

The important contribution a therapeutic relationship makes to supporting, encouraging and motivating patients within the rehabilitation context is generally accepted²¹. Regular social support may well include some of the active elements of this therapeutic dynamic. Psychological support, naturalistic feedback on success or failure of communication, self-monitoring of communication performance, engagement in social activity and networks, demonstration of good conversational models and conversational practice⁹ are also present in social support interventions. Regular provision of such support is likely to increase wellbeing, benefit mood, provide motivation and encourage continued effort amongst a patient group known to be socially isolated as a consequence of their aphasia^{35, 34}.

Regular use of language within a social situation also has the potential to improve functional communication skills through regular conversational practice and naturalistic feedback. Given these benefits, speech and language therapists should ensure that their rehabilitation intervention programme proactively engages and builds the capacity of others in the patient's social circle (volunteers, family members and friends) to provide enhanced social support after stroke related aphasia. This in turn augments the provision of social support beyond what can be provided directly by the therapist. When social support is delivered alongside targeted language rehabilitation it may address the differential dropout and non-adherence rates observed within trials (which suggest that

social support interventions offered as an alternative to therapy maybe unacceptable to people with aphasia). Social support has a role within the rehabilitation and recovery of people with aphasia ⁴ providing functionally relevant opportunities to put into practice gains made during targeted language rehabilitation activities.

From a methodological perspective the distinction between speech and language therapy and social support interventions within these trials has been eroded particularly given that the components of social support are seen as essential elements in speech and language therapy. The distinctions between the interventions were few making it more difficult to establish the effectiveness of those isolated intervention components and increasing the risk of comparison choice bias. Protecting trials from such biases which introduce a consistent tendency for the estimate of effect to differ from the true value is essential to high quality trial design and is particularly challenging when focusing on improving communication. In the context of speech and language therapy trials comparison with an active concurrent control of usual care or another experimental SLT approach would be more acceptable (and is more typical of stroke rehabilitation effectiveness trials in other fields) ⁷.

Active controls in the context of aphasia rehabilitation may differ in the delivery model (professional therapist versus volunteer) ²² or theoretical approach (semantic versus phonological) ²³. Concurrent regimen-controlled interventions might include those that vary in terms of intensity, duration or dose of therapy delivered ^{24, 25}. Control groups may be specifically designed to complement a specific rehabilitation intervention; for example self-directed computerised SLT therapy compared to self-directed computer-based non-language cognitive tasks ^{26, 27}. Such comparisons would ensure that participants across both arms share exposure to the trial processes (i.e. receive the same attention

from the trial staff in assessments, follow-ups, information) but interventions should remain clearly differentiated.

Conclusions:

Interventions that provide social support were used as attention control comparisons in trials of speech and language therapy for aphasia after stroke. The interventions shared numerous similarities and were matched in frequency, intensity, duration, dose and delivery. In several cases social support provider training, support, information, intervention materials and manuals were informed by speech and language therapists. Social support interventions may have been unacceptable to participants with aphasia within trials of therapy effectiveness however as part of a comprehensive language rehabilitation process it is an important adjunct where therapists enable others in the individual's social circle to provide tailored, support and functional language stimulation opportunities for the person with aphasia.

Contributions:

MB: designed the study, reviewed the literature, data extracted, synthesised, analysed and interpreted the data and drafted the manuscript

JG: supported data interpretation and contributed to review of the manuscript

HK: contributed to the interpretation and review of the manuscript

- PE: contributed data, interpreted the data and contributed to drafting and review of the manuscript
- AE: conducted the indirect comparisons, supported interpretation of the data and contributed to drafting and review of the manuscript
- PC: screened full titles, abstracts, full texts, data extracted, interpreted the data and contributed to drafting and review of the manuscript.

CLINICAL MESSAGE

- Social support and speech and language therapy interventions shared many characteristics.
 - Social support is integral to speech and language therapy for aphasia. Therapists work with others to optimise social support approaches, thus extending therapy.
 - More people stopped participating in social support interventions than speech and language therapy making it a poor comparator in evaluations of therapy effectiveness.
-

References

1. Brady MC, Kelly H, Godwin J, Enderby P and Campbell P. Speech and language therapy for aphasia following stroke. *Cochrane Database of Systematic Reviews*. 2016.
2. Braunholtz DA, Edwards SJ and Lilford RJ. Are randomized clinical trials good for us (in the short term)? Evidence for a "trial effect". *J Clin Epidemiol*. 2001; 54: 217-24.
3. Purvis T, Hill K, Kilkenny M, Andrew N and Cadilhac D. Improved in-hospital outcomes and care for patients in stroke research: An observational study. *Neurology*. 2016; 87: 206-13.
4. RCP. National clinical guidelines for stroke. In: Party ISW, (ed.). 5 ed. London 2016.
5. Herbert JD and Gaudiano BA. Moving from empirically supported treatment lists to practice guidelines in psychotherapy: the role of the placebo concept. *J Clin Psychol*. 2005; 61: 893-908.
6. Herbert JD and Gaudiano BA. Introduction to the special issue on the placebo concept in psychotherapy. *J Clin Psychol*. 2005; 61: 787-90.
7. Pollock A, Baer G, Campbell P, et al. Physical rehabilitation approaches for the recovery of function and mobility following stroke. *Cochrane Database Syst Rev*. 2014: CD001920.
8. Higgins JPT and Green S. Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. *The Cochrane Collaboration, 2011 Available from www.cochrane-handbook.org*. 2011.
9. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ*. 2009; 339: b2700.
10. Hoffmann TC, Glasziou PP, Boutron I, et al. Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. *BMJ*. 2014; 348: 1-12.
11. DerSimonian R and Laird N. Meta-analysis in clinical trials. *Controlled Clinical Trials* 1986; 7: 177-88.
12. Bucher HC, Guyatt GH, Griffith LE and SD W. The results of direct and indirect treatment comparisons in meta-analysis of randomized controlled trials. *Journal of clinical epidemiology*. 1997; 50: 683-91.
13. Hedges LV and I. O. *Statistical methods for meta-analysis*. New York: Academic Press, 1985.
14. Woolf C, Cauter A, Haigh Z, et al. A comparison of remote therapy, face to face therapy and an attention control intervention for people with aphasia: a quasi-randomised controlled feasibility study. *Clinical Rehabilitation*. 2016; 30: 359-73.
15. David R, Enderby P and Bainton D. Treatment of acquired aphasia: speech therapists and volunteers compared. *Journal of Neurology, Neurosurgery and Psychiatry*. 1982; 45: 957-61.
16. Lincoln NB, Pickersgill MJ, Hankey AI and Hilton CR. An evaluation of operant training and speech therapy in the language rehabilitation of moderate aphasics. *Behavioural Psychotherapy*. 1982; 10: 162-78.
17. Elman RJ and Bernstein-Ellis E. The efficacy of group communication treatment in adults with chronic aphasia. *Journal of Speech, Language and Hearing Research*. 1999; 42: 411-9.
18. Rochon E, Laird L, Bose A and Scofield J. Mapping therapy for sentence production impairments in nonfluent aphasia. *Neuropsychological Rehabilitation*. 2005; 15: 1-36.
19. Bowen A, Hesketh A, Patchick E, et al. *Clinical Effectiveness, Cost Effectiveness and Service Users' Perceptions of Early, Intensively-Resourced Communication Therapy Following a Stroke, a Randomised Controlled Trial (The ACT NoW Study)*. London: Health Technology Assessment, 2011.
20. Shewan CM and Kertesz A. Effects of speech and language treatment on recovery from aphasia. *Brain and Language*. 1984; 23: 272-99.
21. Wade D. Investigating the effectiveness of rehabilitation professions--a misguided enterprise? *Clin Rehabil*. 2005; 19: 1-3.
22. Meinzer M, Streiftau S and Rockstroh B. Intensive language training in the rehabilitation of chronic aphasia - effective training by laypersons. *Journal of the International Neuropsychological Society*. 2007; 13: 846-53.

23. Doesborgh SJC, Van de Sandt-Koenderman MWE, Dippel DWJ, Koudstaal PJ and Visch-Brink EG. Effects of semantic treatment on verbal communication and linguistic processing in aphasia after stroke: a randomized controlled trial. *Stroke*. 2004; 35: 141-6.
24. Martins IP, Leal G, Fonseca I, et al. A randomized, rater-blinded, parallel trial of intensive speech therapy in sub-acute post-stroke aphasia: the SP-I-R-IT study. *Int J Lang Commun Disord*. 2013; 48: 421-31.
25. Godecke E, Hird K, Lalor EE, Rai T and Phillips MR. Very early poststroke aphasia therapy: a pilot randomized controlled efficacy trial. *International Journal of Stroke*. 2012; 7: 635-44.
26. Katz RC and Wertz RT. The efficacy of computer-provided reading treatment of chronic aphasic adults. *Journal of Speech, Language and Hearing Research*. 1997; 40: 493-507.
27. Palmer R, Cooper C, Enderby P, et al. Clinical and cost effectiveness of computer treatment for aphasia post stroke (Big CACTUS): study protocol for a randomised controlled trial. *Trials*. 2015; 16: 18-.
28. Bhogal SK, Teasell R and Speechley M. Intensity of aphasia therapy, impact on recovery. *Stroke*. 2003; 34: 987-93.
29. Robey R. The efficacy of treatment for aphasic persons: a meta-analysis. *Brain and Language*. 1994; 47: 582-608.
30. Robey R. A meta-analysis of clinical outcomes in the treatment of aphasia. *Journal of Speech, Language and Hearing Research*. 1998; 41: 172-87.
31. Cherney LR, Patterson JP, Raymer A, Frymark T and Schooling T. Evidence-based systematic review: effects of intensity of treatment and constraint-induced language therapy for individuals with stroke-induced aphasia. *Journal of Speech, Language and Hearing Research*. 2008; 51: 1282-99.
32. Shewan CM, Bandur DL. Treatment of Aphasia: A Language-oriented Approach. San-Diego: College-Hill Press, 1986.
33. Bell ML, Kenward MG, Fairclough DL, Horton NJ. Differential dropout and bias in randomised controlled trials: when it matters and when it may not *BMJ* 2013; 346 :e8668
34. Hilari K, Northcott S. "Struggling to stay connected": comparing the social relationships of healthy older people and people with stroke and aphasia, *Aphasiology* 2016; 31:6, 674-687, DOI: 10.1080/02687038.2016.1218436
35. Parr S. Living with severe aphasia: Tracking social exclusion, *Aphasiology*, 2011: 21(1)98-123, DOI: 10.1080/02687030600798337
36. Schulz KF, Altman DG, Moher D, for the CONSORT Group. CONSORT 2010 Statement: updated guidelines for reporting parallel group randomised trials. *BMC Medicine* 2010, 8:18. (24 March 2010)
37. Skehon M, Cartwright M, Francis JJ. Acceptability of healthcare interventions: an overview of reviews and development of a theoretical framework. *BMC Health Services Research* 2017**17**:88. doi.org/10.1186/s12913-017-2031-8